

Examination of Apoptotic Effects of High-Dose Toluene on the Brain Cortex and Cerebellum Tissue During the Acute Phase: an experimental study

Ufuk Tas¹, Murat Ayan², Tuncay Kuloglu³, Mustafa Suren⁴, Duygu Cakıl⁵, Birsen Ozyurt¹, Dürdane Aksoy Bekar⁶, Mustafa Sarsılmaz⁷

ABSTRACT

The aim of this study is to examine the apoptotic effects of high-dose toluene on rat brain cortex and cerebellum tissues during the acute phase. Fourteen albino Wistar rats (180-200 g) were divided into two equal groups (n : 7). While Group I was used as a control group, Group II was applied to a high dose of toluene (5200 mg / kg / gavage). At the end of a three-hour experimental period brain tissue samples were taken from the killed animals and tissues were fixed in % 10 neutral formalin, then, embedded in paraffin and sectioned (thickness, 5 µm) to determine the immune reactivity of Bax, sections were stained immunohistochemically with avidin biotin-peroxidase method. At the same time the TUNEL method was used for detection of apoptosis in the brain cortex and cerebellum. As a result of the study, increased Bax immune reactivity was seen in the brain cortex (+++) and cerebellum (++++) of toluene treated rats compared to control. The numbers of TUNEL-positive cells were significantly higher in the cerebellum tissues of animals exposed to toluene than the control. Nevertheless there was not a significantly difference in terms of TUNEL positivity between the brain cortex of control and toluene exposed animals. As a result of this study it was shown that a high-dose of toluene can trigger apoptosis of the brain cortex and cerebellum in a very short period of time.

Key words: Toluene, apoptosis, brain cortex and cerebellum, rat.

Yüksek Doz Tolüenin Akut Dönemde Beyin Korteksi ve Serebellum Dokusu Üzerindeki Apoptotik Etkilerinin İncelenmesi; Deneysel Bir Çalışma

ÖZET

Bu çalışmada yüksek doz tolüenin rat beyin korteksi ve serebellum dokuları üzerinde akut dönemdeki apoptotik etkilerinin araştırılması amaçlandı. On dört adet Wistar Albino cinsi sıçan (180-200 g) iki eşit guruba (n:7) bölündü. Grup I kontrol grubu olarak kullanılırken Grup II'ye yüksek doz tolüen verildi (5200 mg/kg/gavaj). Üç saatlik deney periyodunun sonunda öldürülen hayvanlardan beyin doku örnekleri % 10'luk nötral formaline alınıp parafine gömülerek kesildi (5 µm). Bax immün reaktivitesinin belirlenebilmesi için kesitler avidin biotin peroksidaz yöntemi ile immünohistokimyasal olarak boyandı. Aynı zamanda beyin korteksi ve serebellumda apoptozun tespit edilmesi için TUNEL yöntemi uygulandı. Çalışma sonucunda tolüen verilen sıçanlarda beyin korteksi (+++) ve serebellumda (++++) artmış Bax immün reaktivitesine rastlanıldı. Tolüene maruz kalan hayvanların serebellum dokularında TUNEL pozitif hücre sayısı kontrol grubuna oranla anlamlı oranda yüksekti. Bununla birlikte beyin korteksi dokusunda tolüen ve kontrol grupları arasında TUNEL pozitifliği açısından anlamlı bir fark yoktu. Bu çalışma sonucunda yüksek doz tolüenin beyin korteksi ve serebellumda çok kısa sürede apoptozu tetikleyebileceği gösterilmiştir.

Anahtar kelimeler: Tolüen, apoptozis, beyin korteksi ve serebellum, sıçan

Gaziosmanpaşa University, Faculty of Medicine, Departments of Anatomy¹, Emergency Medicine², Anesthesiology and Reanimation¹, Physiology³ and Neurology⁴ Tokat, ⁵Firat University, Faculty of Medicine, Department of Histology and Embryology, Elazığ, ⁶Sifa University, Faculty of Medicine Department of Anatomy, Izmir, Turkey

Correspondence: Ufuk Taş, MD
Gaziosmanpaşa University, Faculty of Medicine, Department of Anatomy,
60000 Tokat, Turkey.
Tel: +90 356 2129500 (1232)
Fax: +90 356 2129417
E-mail: dr_ufuktas@hotmail.com

Received: 26.04.2012, Accepted: 28.04.2012

INTRODUCTION

Toluene is a liquid aromatic hydrocarbon that widely used in industry and addiction is observed commonly (1-3). The abuse of toluene like volatile substances is an important problem in the many countries, using of this substance with up to 10% to 15% of young people developed country (4). Due to the high percentage of toluene, addicts mostly prefer sniffing glue and paint thinner (2, 5, 6). Toluene can be taken through the respiratory, digestive system and the skin into the body and generally accumulates at highly vascularized tissues which are rich in terms of fat, a large part is destroyed in the liver and converted into hippuric acid and then is excreted through urine (1, 7, 8). Toluene may cause many clinical situations from a headache up to death depending on the dose and duration (1, 9). Exposure to high-dose toluene generally is observed at workers and dependents (10, 11). Researches on high-dose poisonings of toluene are mostly case studies based on the evaluation of cases of acute intoxication (12, 13). Since toluene mainly accumulates in nervous tissue, mostly chronic effects of toluene on the nervous system were investigated. Although, there are many studies examining the effects of chronic exposure to brain tissue (4, 14-16), only limited experimental studies examining the damage of acute and high doses of toluene (17, 18). Furthermore we didn't find any study about toluene induced apoptosis in brain and cerebellum tissue within a few hours. In this study we aimed to investigate the apoptotic effects of high-dose toluene on rat brain cortex and cerebellum tissues during the acute phase using immunohistochemical methods.

MATERIAL AND METHODS

Animals

Fourteen adult male Wistar-albino rats (250-300 g) obtained from Gaziosmanpasa University Biomedical Research Unit were randomly divided into two equal groups: Group I (n:7) as control and Group II (n:7) as treated with toluene. The experimental protocols were approved (2011 HADYEK-048) by the appropriate animal care committee of Gaziosmanpasa University.

Chemicals, route of exposure and dose selection

Toluene was administered by gavage without dilution described in previous study (19). The acute oral LD50 of toluene in adult rats ranged from 5.5 to 7.4 g/kg

Table 1. The scoring of the immune reactivity (Bax)

Grade	Symbol
absent	(-)
very few	(±)
few	(+)
medium	(++)
high	(+++)
very high	(++++)

(20). We defined the benchmark dose as the maximum neurotoxic dose causing toxic damage without leading to death (19). Rats in Group I were administered serum through gavage, while Group II rats were given a single dose (5200 mg/kg/gavage) of 99.5% pure toluene (Sigma, St. Louis, Missouri, USA). Sample collection and preparations of brain cortex and cerebellum tissue. At the end of the three-hour experimental period all rats were killed by exsanguination under ketamine/xylazine (50/10 mg/kg) anesthesia. Brain tissues were removed directly and fixed in formalin solution for immunohistochemical evaluations. The paraffin-embedded brain specimens were cut into 4-5 µm sections and stained with Bax for immunohistochemical evaluation and TUNEL for apoptosis determination. Specimens were examined under a Novel N-800M light microscope.

Immunohistochemistry

Paraffin embedded brain cortex and cerebellum tissues were dissected at 4-5 µm and deparaffinized in xylene, then dehydrated with alcohol series. The brain cortex and cerebellum tissue were then placed in distilled water and boiled in citrate buffer solution (pH:6.0) in a microwave oven (750W) for 7+5 minutes for antigen retrieval. To prevent endogenous peroxidase activity sections were treated with 3% hydrogen peroxide.

Table 2. Semi-quantitative evaluation of Bax staining in brain cortex and cerebellum tissues of control and toluene-treated rats (n: 7 for each group).

Tissue	Control	Toluene
Brain cortex	(+)	(+++)
Cerebellum	(+)	(++++)

The number of the positive staining was recorded as very few (±), few (+), medium (++) , high (+++) and very high (++++).




Figure 1. Bax staining in the brain cortex (400x) and cerebellum (200x) tissues of the animals in the toluene and control group. A small amount of BAX (+) positive cells in the brain cortex (A) and cerebellum (D) of the control group. Negative control of the brain cortex (B) and cerebellum (E) tissue of animals exposed to toluene. BAX positive cells in the brain cortex (C) (+++), and cerebellum (F) (++++) tissues of animals exposed to toluene.

Tissues were treated with Ultra V Block (Ultra V Block, TA-125-UB, Thermo Fisher Scientific Inc, USA) solution to prevent background staining and then incubated with primer antibody Bax (mouse monoclonal IgG, Santa Cruz Biotechnology, sc-32239, California, USA) for 60 minutes. Secondary antibody application (biotinated anti-mouse IgG, Diagnostic BioSystems, KP 50A, Pleasanton, USA) was performed for 30 minutes. After streptavidin horseradish peroxidase treatment for 30 minutes and 3-amino-9-ethyl carbazole chromogen treatment, contrast staining was carried out using Mayer's hematoxylin. Phosphate buffered saline (PBS) was used instead of primary antibody for the negative control. Tissues treated with PBS and distilled water were covered with an appropriate covering solution.

TUNEL assay

ApopTag plus Peroxidase in Situ Apoptosis Detection Kit (Chemicon, Cat no: S7101, USA) were used for detection of apoptotic cells. According to the instructions of the manufacturer sections were deparaffinized in xylene, dehydrated through graded alcohol, and washed in PBS. Tissues were incubated in a 0.05% proteinase K solution and were incubated with 3% hydrogen peroxide for five minutes to prevent endogenous peroxidase activity. Later washing with PBS, the tissues were placed in equilibration buffer for six minutes and in working solu-




Figure 2. TUNEL staining in the brain cortex (400x) and cerebellum (200x) tissues of the animals in the toluene and control group. The brain cortex (A) and cerebellum (B) tissues of the control group. While in the brain cortex (C) there was no significant difference between toluene group and control group in terms of apoptotic cells, a significantly number of apoptotic cells in the cerebellum tissue (D) are observed. Negative controls of the brain cortex (E) and cerebellum (F) tissues of animals exposed to toluene. Positive control staining in the gastric tissue (G).

tion (70% reaction buffer plus 30% TdT enzyme) at 37°C under moist conditions for one hour. Stop/wash buffer were applied during for 10 minutes and then anti-digoxigenin-peroxidase for 30 minutes. Diaminobenzidine (DAB) substrate was used to showing apoptotic cells. Sections were counterstained with methyl green. Stomach tissue was used for a positive control. PBS was used instead of the Tdt enzyme on the negative control. Cells with green nuclei after TUNEL staining using methyl green were considered normal, whereas cells with brown nuclei were considered apoptotic.

Evaluations of tissue sections

Preparations were observed and photographed (Figure 1 and Figure 2) using a research microscope (Novel N-800M). Apoptotic (TUNEL positive) cells were counted in at least eight areas per tissue section, in two sections from each animal, at 400X magnification. Bax staining was evaluated according to the method described previously (21) (Table 1).




Figure 3. Apoptotic index of TUNEL positive cell in the brain cortex and cerebellum tissues of the animals in the toluene and control group a, $p < 0.01$ compared to control group in cerebellum a, $p > 0.05$ compared to control group in brain cortex

Statistical analyses

Results are expressed as mean \pm standard deviation (SD). Mann-Whitney U test were used for statistical analysis. All analyses were performed using SPSS version 15.0 software. P values of < 0.05 were considered to be statistically significant.

RESULTS

Immunohistochemical findings

Bax protein was visualized by immunohistochemical staining of brain cortex and cerebellum tissues cross-sections of rats in the toluene and control groups (Figure 1). The results were evaluated semi-quantitatively. There were significant increases in Bax immune reactivity in the brain cortex (+++) and cerebellum (++++), tissue of rats in the toluene group, whereas there were little staining (+) in the control group (Table 2).

TUNEL findings

TUNEL staining was shown in the brain cortex and cerebellum tissue sections of rats in the toluene and the control group (Figure 2). The numbers of TUNEL-positive cells in the cerebellum tissues of animals exposed to toluene were significantly higher than the control group ($p < 0.05$). However there was not a significantly difference in terms of TUNEL positivity between the brain cortex of control and toluene exposed animals (Figure 3).

DISCUSSION

The majority of toluene is broken down via the cytochrome P-450 enzyme found in the liver and converted to hippuric acid, the major and last metabolite of toluene, which is excreted in the urine (22). Toluene and its metabolites affect the defense system in the body by means of oxidative stress and may cause damage in many tissues primarily including the brain (16, 23-25). While the mechanism of action of toluene is not exactly clear, authors mention several different mechanisms. Toluene modifies lipid composition of cell membrane and effects the Na / K-ATPase activity and thus increases the membrane fluidity (26). Some researchers have reported that toluene may affect Gabaergic, glutamatergic, serotonergic, dopaminergic pathways (2, 27). In addition, it is said that toluene may cause mitochondrial damage, and increase of pro-apoptotic proteins (Bax) in the cytoplasm and thus increases apoptotic activity (28, 29). Apoptosis known as controlled cell death without causing inflammation, it is characterized by cytoplasm shrinkage, membrane changes, and DNA fragmentation. Recently it is mentioned that apoptosis may be triggered with various factors and follow different pathways (30). In mitochondria-mediated apoptosis, which is the most popular of these, apoptosis is triggered by releasing of apoptogenic factors like, cytochrome c, apoptosis-inducing factor and smac / DIABLO from intermembranal space of mitochondria to cytosol. Cytochrome C released into the cytosol, the apoptosis complex consisting of C/Apaf-1/ATP/proxaptase-9 first activates caspase-9 and then caspase-3 (31, 32). Family member of Bcl-2 are effective in the increase or decrease of apoptotic activity. During the apoptosis period pro-apoptotic mediators which passing from the mitochondria into the cytosole (Bax, Bak, Bad), accelerate the apoptotic activity by increasing the release of cytochrome c (33, 34). Chromatin condensation, DNA fragmentation and catabolism of membrane proteins like morphological changes occur with Caspase-3 activation. (35). During apoptosis, these protein changes can be determined by immunohistochemical staining method and DNA damage can be determined with TUNEL staining method. In our study significantly increased Bax-positive cells was seen in the brain cortex and cerebellum tissues of rats exposed to toluene. This increase was evaluated in favor of the increase in apoptotic activity. The number of TUNEL-positive cells in the cerebellum tissues of animals exposed to toluene was significantly higher than

in the control group. However no significant difference was seen between the toluene group and control group in terms of TUNEL positivity in the brain cortex. El-Nabi Kamel and Shehata (2008) reported that the most affected tissue by toluene is the brain tissue (23). It was shown that toluene causes damage to the frontal cortex and brain stem and leads to an increase in apoptotic activity (36). Baydas et al have shown morphological changes (GFAP) in the cerebellum, cortex and hippocampus tissues of 60-70% toluene containing thinner exposed animals (37). Gotohda et al were demonstrated shrunken granule cells in the dentate gyrus of the hippocampus and shrunken Purkinje cells in cerebellum after toluene inhalation in rat brain (15). Hester et al demonstrated global gene expression changes in rat brain after toluene inhalation during an acute exposure. They found toluene exposure was associated with changing of genes in pathways associated with GABA receptor signaling and mitochondrial function (18). Heung-Sik Seo et al claim that acute toluene exposure (0-1000 mg/kg) reduces the rate of adult hippocampal neurogenesis but does not induce neural apoptosis in the hippocampus of adult mice (17). Kanter was demonstrated the significantly increased number of apoptotic neurons in chronic toluene treated rats compared to control animals in frontal cortex tissues using by TUNEL assay (14). Considering previous studies, it can be said that as a result of chronic toluene exposure especially may cause apoptotic changes and structural damage in the brain cortex and cerebellum tissue. Although the number of experimental studies in relation to acute exposure is very limited, it is mentioned some of these studies that toluene stimulates apoptosis lightly or does not induce apoptosis. However in our study, significantly increased Bax immunoreactivity was shown in the brain cortex and cerebellum tissues of animals given high doses of toluene, and that toluene is able to trigger apoptosis in a very short time like 3 hours. The numbers of apoptotic cells (TUNEL positive cell) in the cerebellum tissues of toluene treated animals were significantly higher but not in cortex compared to control. It may be concluded from this study, cortex of the brain tissue is affected more quickly than cerebellum in toluene poisoning. For better understanding of the pathogenesis of acute intoxication cases, it is necessary to make further studies in relation to acute high-dose exposure to toluene on the molecular level in the future.

REFERENCES

1. Agency for Toxic Substances and Disease Registry (ATSDR) (2006) *Case Studies in Environmental Medicine; Toluene Toxicity* <http://www.atsdr.cdc.gov/csem/toluene/docs/toluene.pdf> (accessed october 10, 2011).
2. Balster RL. Neural basis of inhalant abuse. *Drug Alcohol Depend* 1998;51:207-14.
3. Jones HE, Balster RL. Neurobehavioral consequences of intermittent prenatal exposure to high concentrations of toluene. *Neurotoxicol Teratol* 1997;19:305-13.
4. Filley CM, Halliday W, Kleinschmidt-DeMasters BK. The effects of toluene on the central nervous system. *J Neuropathol Exp Neurol* 2004;63:1-12.
5. Beyer CE, Stafford D, LeSage MG, Glowa JR, Steketee JD. Repeated exposure to inhaled toluene induces behavioral and neurochemical cross-sensitization to cocaine in rats. *Psychopharmacology (Berl)* 2001;154:198-204.
6. Park CK, Kwon KT, Lee DS, et al. [A case of toxic hepatitis induced by habitual glue sniffing]. *Taehan Kan Hakhoe Chi* 2003 ;9:332-6.
7. Faust RA. Chemical summary for toluene. 1994. http://www.epa.gov/chemfact/s_toluen.txt (accessed march 21, 2012).
8. Dossing M, Aelum JB, Hansen SH, Lundqvist GR, Andersen NT. Urinary hippuric acid and orthocresol excretion in man during experimental exposure to toluene. *Br J Ind Med* 1983;40:470-3.
9. OEHHA, Office of Environmental Health Hazard Assessment. 2004. *Public health goals for chemicals in drinking water*. <http://www.oehha.org/water/phg/pdf/perchlorate3docs.pdf> (accessed January 21, 2012).
10. Tsatsakis AM, Dolapsakis G, Troulakis G, Christodoulou P, Relakis K, Trikilis N, et al. Fatal and non-fatal outcome by accidental intoxication with paint thinner. *J Clin Forensic Med* 1997;4:133-7.
11. Tomei F, Giuntoli P, Biagi M, Baccolo TP, Tomao E, Rosati MV. Liver damage among shoe repairers. *Am J Ind Med* 1999;36:541-7.
12. Argo A, Bongiorno D, Bonifacio A, Pernice V, Liotta R, Indelicato S, et al. A fatal case of a paint thinner ingestion: comparison between toxicological and histological findings. *Am J Forensic Med Pathol* 2011;31:186-91.
13. Tan WP, Seow E. Case reports on acute toluene poisoning during parquet flooring. *Ann Acad Med Singapore* 1997;26:138-40.
14. Kanter M. Protective effects of thymoquinone on the neuronal injury in frontal cortex after chronic toluene exposure. *J Mol Histol* 2011;42:39-46.
15. Gotohda T, Tokunaga I, Kubo S, Morita K, Kitamura O, Eguchi A. Effect of toluene inhalation on astrocytes and neurotrophic factor in rat brain. *Forensic Sci Int* 2000;113:233-8.
16. Baydas G, Ozveren F, Akdemir I, Tuzcu M, Yasar A. Learning and memory deficits in rats induced by chronic thinner exposure are reversed by melatonin. *J Pineal Res* 2005;39:50-6.

17. Seo HS, Yang M, Song MS, Kim JS, Kim SH, Kim JC, et al. Toluene inhibits hippocampal neurogenesis in adult mice. *Pharmacol Biochem Behav* 2010;94:588-94.
18. Hester SD, Johnstone AF, Boyes WK, Bushnell PJ, Shafer TJ. Acute toluene exposure alters expression of genes in the central nervous system associated with synaptic structure and function. *Neurotoxicol Teratol* 2011;33:521-9.
19. Mehta CS, Sun PS, Zikarge A. Acute Toxicity of Toluene In Male and Female Rats: A Single Oral Dose Exposure 2-Week Study. *Toxic Substance Mechanisms* 1998;17:43-56.
20. Agency for Toxic Substances and Disease Registry (ATSDR) (2000) Toxicological Profile for Toluene, <http://www.atsdr.cdc.gov/toxprofiles/tp56.pdf> (accessed october 15, 2011).
21. Kanter M. Thymoquinone attenuates lung injury induced by chronic toluene exposure in rats. *Toxicol Ind Health* 2011;27:387-95.
22. Pero RW. Health consequences of catabolic synthesis of hippuric acid in humans. *Curr Clin Pharmacol* 2010;5:67-73.
23. El-Nabi Kamel MA, Shehata M. Effect of toluene exposure on the antioxidant status and apoptotic pathway in organs of the rat. *Br J Biomed Sci* 2008;65:75-9.
24. Gotohda T, Nishimura A, Morita K. Immunohistochemical studies on early stage of hepatic damage induced by sub-acute inhalation of toluene vapor in rats. *J Appl Toxicol* 2009;29:505-9.
25. Kanter M. Thymoquinone reestablishes spermatogenesis after testicular injury caused by chronic toluene exposure in rats. *Toxicol Ind Health* 2011;27:155-66.
26. Calderon-Guzman D, Espitia-Vazquez I, Lopez-Dominguez A, Hernandez-Garcia E, Huerta-Gertrudis B, Coballase-Urritia E, et al. Effect of toluene and nutritional status on serotonin, lipid peroxidation levels and $NA^+/K^+-ATPase$ in adult rat brain. *Neurochem Res* 2005;30:619-24.
27. Bowen SE, Batis JC, Mohammadi MH, Hannigan JH. Abuse pattern of gestational toluene exposure and early postnatal development in rats. *Neurotoxicol Teratol* 2005;27:105-16.
28. Lee JI, Lee KS, Paik YH, Nyun Park Y, Han KH, Chon CY, et al. Apoptosis of hepatic stellate cells in carbon tetrachloride induced acute liver injury of the rat: analysis of isolated hepatic stellate cells. *J Hepatol* 2003;39:960-6.
29. Tas U, Ogeturk M, Meydan S, Kus I, Kuloglu T, Ilhan N, et al. Hepatotoxic activity of toluene inhalation and protective role of melatonin. *Toxicol Ind Health* 2011;27:465-73.
30. Saraste A, Pulkki K. Morphologic and biochemical hallmarks of apoptosis. *Cardiovasc Res* 2000;45:528-37.
31. Bender LM, Morgan MJ, Thomas LR, Liu ZG, Thorburn A. The adaptor protein TRADD activates distinct mechanisms of apoptosis from the nucleus and the cytoplasm. *Cell Death Differ* 2005;12:473-81.
32. Fulda S, Debatin KM. Resveratrol modulation of signal transduction in apoptosis and cell survival: a mini-review. *Cancer Detect Prev* 2006;30:217-23.
33. Win-Shwe TT, Yoshida Y, Kunugita N, Tsukahara S, Fujimaki H. Does early life toluene exposure alter the expression of NMDA receptor subunits and signal transduction pathway in infant mouse hippocampus? *Neurotoxicology* 2010;31:647-53.
34. Tao XJ, Tilly KI, Maravei DV, Shifren JL, Krajewski S, Reed JC, et al. Differential expression of members of the bcl-2 gene family in proliferative and secretory human endometrium: Glandular epithelial cell apoptosis is associated with increased expression of bax. *Journal of Clinical Endocrinology & Metabolism* 1997;82:2738-46.
35. D'Amelio M, Cavallucci V, Cecconi F. Neuronal caspase-3 signaling: not only cell death. *Cell Death Differ* 2010;17:1104-14.
36. Kanter M. Protective effects of *Nigella sativa* on the neuronal injury in frontal cortex and brain stem after chronic toluene exposure. *Neurochem Res* 2008;33:2241-9.
37. Baydas G, Reiter RJ, Nedzvetskii VS, Yasar A, Tuzcu M, Ozveren F, et al. Melatonin protects the central nervous system of rats against toluene-containing thinner intoxication by reducing reactive gliosis. *Toxicology Letters* 2003;137:169-74.